

| AAHA Vaccine Protocols for Canines | | | |
|--|---|--|--|
| | Primary vacc <16 wks | Primary vacc >16 wks | Booster |
| CORE | | | |
| Canine distemper virus (MLV) | 6-8 weeks, then 9-11 weeks, then 12-14 weeks | 1 dose adequate (2 doses 3-4 weeks apart is acceptable) | first 1 year, then every 3 years |
| Parvovirus (MLV) | 6-8 weeks, then 9-11 weeks, then 12-14 weeks | 1 dose adequate (2 doses 3-4 weeks apart is acceptable) | first 1 year, then every 3 years |
| Canine adenovirus-2 (MLV or killed) | 6-8 weeks, then 9-11 weeks, then 12-14 weeks | MLV: 1 dose adequate (2 doses 3-4 weeks apart is acceptable) killed: 2 doses 2-4 weeks apart | first 1 year, then every 3 years |
| Rabies virus (killed; 3 year label) | 1 dose as early as 3 months (or as per state law) | 1 dose | first 1 year, then every 3 years (or as per state law) |
| NON-CORE | | | |
| Canine parainfluenza virus (MLV) | 6-8 weeks, then 9-11 weeks, then 12-14 weeks | 1 dose adequate (2 doses 3-4 weeks apart is acceptable) | first 1 year, then every 3 years |
| Bordetella (live avirulent) + CPiV (MLV) + CAV-2 (MLV) (intranasal) | 1 dose (start no younger than 8 weeks) | 1 dose | annual (if not vaccinated within last 6 months, booster 1 week prior to known exposure, such as boarding) |
| Lyme (killed whole bacterin or recombinant OspA) | at least 9 weeks (optimally at least 3 months), then 3-4 weeks later | 2 doses 2-4 weeks apart | annual |
| Leptospira (killed bacterin) | 12 weeks, then 16 weeks (start no younger than 12 weeks) | 2 doses 2-4 weeks apart | annual (if risk of exposure to wildlife, livestock, or waterways*; every 6 months if <i>significant</i> risk) |
| NOT RECOMMENDED | | | |
| Coronavirus, Giardia, Canine adenovirus-1 | | | |

*based on recommendation of UCD VMTH Outpatient Service, Dr. Gina Davis